

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

### **Supplemental Material**

#### **Evaluating the Effect of Gestational Exposure to Perfluorohexane Sulfonate on Placental Development in Mice Combining Alternative Splicing and Gene Expression Analyses**

Yihao Zhang, Jia Lv, Yi-Jun Fan, Lin Tao, Jingjing Xu, Weitian Tang, Nan Sun, Ling-Li Zhao, De-Xiang Xu, and Yichao Huang

#### **Table of Contents**

**Figure S1.** Experimental animal phenotype. PFHxS exposure from GD7 to GD17 for pregnant mice (mean  $\pm$  SEM). The exposure dose were 0.3 (High) and 0.03 (Low)  $\mu\text{g/kg/day}$ , which were administered in corn oil with 0.1% DMSO; and the Control was corn oil with 0.1% DMSO. (A) Maternal weight; (B) Food consumption; (C) Liver/Body weight ratio; (D) Brain/Body weight ratio; (E) Liver births per litter.  $N = 6$  in each group,  $P$  values were determined by One-way Analysis of Variance, all  $P > 0.05$ . Data in (A-E) are also presented in Excel Table S18. Notes: DMSO, dimethyl sulfoxide; GD, gestational day; PFHxS, perfluorohexane sulfonate; SEM, Standard Error of Mean.

**Figure S2.** The distribution of core genes in GSEA. (A) The distribution of core genes in GSEA for placenta development. (B) The distribution of core genes in GSEA for multicellular organism metabolic process. (C) The distribution of core genes in GSEA for anion transmembrane transport.  $N = 6$  in each group, including 3 males and 3 females.  $P$  and FDR values were determined by permutation test and Benjamini-Hochberg method, respectively. Data in (A-C) are also presented in Excel Table S19. Notes: FDR, false discovery rate; GO, Gene Ontology; GSEA, gene set enrichment analysis; PFHxS, perfluorohexane sulfonate.


**Figure S3.** Signal pathways in multiple biological processes of placental development in response to PFHxS exposure. (A) Heatmap showing the DEGs profile of development gene set between high-dose group and control. (B) Heatmap showing the DEGs profile of metabolic process gene set between high-dose group and control. (C) Heatmap showing the DEGs profile of proliferation gene set between high-dose group and control. (D) Heatmap showing the DEGs profile of transport gene set between high-dose group and control. (E) GSEA showing the positive enrichment of TGF  $\beta$  signaling pathway in high-dose group compare to control. (F) GSEA showing the positive enrichment of p53 signaling pathway in high-dose group compare to control. (G) GSEA showing the positive enrichment of Wnt signaling pathway in high-dose group compare to control. (H) UpSetR plot showing the number of unique and shared DEGs between development, proliferation, metabolic process and transport gene sets. (I) Circos plot showing GO analysis of shared DEGs for 4 gene sets.  $N = 6$  in each group, including 3 males and 3 females.  $P$  and FDR values were determined by permutation test and Benjamini-Hochberg method (E, F and G), respectively. Data in (A-I) are also presented in Excel Table S20. Notes: DEGs, differentially expressed genes; FDR, false discovery rate; GSEA, gene set enrichment analysis; PFHxS, perfluorohexane sulfonate.

**Figure S4.** AS analyses in multiple signal pathways of metabolism and transport after PFHxS exposure. (A) The number of differential AS events in male placenta with PFHxS expose compare to control. (B) The number of differential AS events in female placenta with PFHxS expose compare to control. (C) The proportion of differential AS events in male and female placenta with PFHxS expose compare to control. (D) GO analysis showing the genes with AS event are enriched in multiple signal pathways of metabolism and transport in male placenta. (E) GO analysis showing the genes with AS event are enriched in multiple signal pathways of metabolism and transport in female placenta. (F) Heatmap showing the profile of splicing factors in male and female placenta with PFHxS expose compare to control.  $N = 6$  in each group, including 3 males and 3 females.  $P$  values were determined by hypergeometric test (D and E). Data in (A-F) are also presented in Excel Table S21. Notes: AS, alternative splicing; GO, Gene Ontology; PFHxS, perfluorohexane sulfonate.


**Figure S5.** AS analyses in multiple signal pathways of placental development after PFHxS exposure. (A) KEGG pathway interaction analysis showing the relationship between signal pathways in RI gene set. (B) KEGG pathway interaction analysis showing the relationship between signal pathways in A3SS gene set. (C) KEGG pathway interaction analysis showing the relationship between signal pathways in A5SS gene set. (D) KEGG pathway interaction analysis showing the relationship between signal pathways in SE gene set. (E) KEGG pathway interaction analysis showing the relationship between signal pathways in MXE gene set. (F) Reactome analysis showing the genes with AS event are enriched in multiple signal pathways related to placental development.  $N = 6$  in each group, including 3 males and 3 females.  $P$  values were determined by hypergeometric test (F). Data in (A-F) are also presented in Excel Table S22. Notes: A3SS, alternative 3' splice site; A5SS, alternative 5' splice site; AS, alternative splicing; KEGG, Kyoto Encyclopedia of Genes and Genomes; MXE, exclusion exon; SE skipped exon; PFHxS, perfluorohexane sulfonate; RI, retained intron.

**Figure S6.** Evaluation of placental amino acid transport in the high-dose group and control in both genders. (A) The quantitative analysis of mRNA of 12 amino acid transporters in placenta ( $N = 6$ , mean  $\pm$  SEM). (B) The profile of 29 amino acids and 44 amides in male placenta ( $N = 5$ ). (C) The profile of 29 amino acids and 44 amides in female placenta ( $N = 5$ ).  $P$  values were determined by Student's  $t$  test (A),  $*P < 0.05$ . Data in (A-C) are also presented in Excel Table S23. Note: PFHxS, perfluorohexane sulfonate; SEM, Standard Error of Mean.

**Additional File-** Excel Document




**Figure S1.** Experimental animal phenotype. PFHxS exposure from GD7 to GD17 for pregnant mice (mean  $\pm$  SEM). The exposure dose were 0.3 (High) and 0.03 (Low)  $\mu\text{g/kg/day}$ , which were administered in corn oil with 0.1% DMSO; and the Control was corn oil with 0.1% DMSO. (A) Maternal weight; (B) Food consumption; (C) Liver/Body weight ratio; (D) Brain/Body weight ratio; (E) Live births per litter.  $N = 6$  in each group,  $P$  values were determined by One-way Analysis of Variance, all  $P > 0.05$ . Data in (A-E) are also presented in **Excel Table S18**. Notes: DMSO, dimethyl sulfoxide; GD, gestational day; PFHxS, perfluorohexane sulfonate; SEM, Standard Error of Mean.





**Figure S2.** The distribution of core genes in GSEA. **(A)** The distribution of core genes in GSEA for placenta development. **(B)** The distribution of core genes in GSEA for multicellular organism metabolic process. **(C)** The distribution of core genes in GSEA for anion transmembrane transport.  $N = 6$  in each group, including 3 males and 3 females.  $P$  and FDR values were determined by permutation test and Benjamini-Hochberg method, respectively. Data in **(A-C)** are also presented in **Excel Table S19**. Notes: FDR, false discovery rate; GO, Gene Ontology; GSEA, gene set enrichment analysis; PFHxS, perfluorohexane sulfonate.

**A**





**B**


















**Figure S3.** Signal pathways in multiple biological processes of placental development in response to PFHxS exposure. (A) Heatmap showing the DEGs profile of development gene set between high-dose group and control. (B) Heatmap showing the DEGs profile of metabolic process gene set between high-dose group and control. (C) Heatmap showing the DEGs profile of proliferation gene set between high-dose group and control. (D) Heatmap showing the DEGs profile of transport gene set between high-dose group and control. (E) GSEA showing the positive enrichment of TGF  $\beta$  signaling pathway in high-dose group compare to control. (F) GSEA showing the positive enrichment of p53 signaling pathway in high-dose group compare to control. (G) GSEA showing the positive enrichment of Wnt signaling pathway in high-dose group compare to control. (H) UpSetR plot showing the number of unique and shared DEGs between development, proliferation, metabolic process and transport gene sets. (I) Circos plot showing GO analysis of shared DEGs for 4 gene sets.  $N = 6$  in each group, including 3 males and 3 females.  $P$  and FDR values were determined by permutation test and Benjamini-Hochberg method (E, F and G), respectively. Data in (A-I) are also presented in **Excel Table S20**. Notes: DEGs, differentially expressed genes; FDR, false discovery rate; GSEA, gene set enrichment analysis; PFHxS, perfluorohexane sulfonate.




**Figure S4.** AS analyses in multiple signal pathways of metabolism and transport after PFHxS exposure. (A) The number of differential AS events in male placenta with PFHxS expose compare to control. (B) The number of differential AS events in female placenta with PFHxS expose compare to control. (C) The proportion of differential AS events in male and female placenta with PFHxS expose compare to control. (D) GO analysis showing the genes with AS event are enriched in multiple signal pathways of metabolism and transport in male placenta. (E) GO analysis showing the genes with AS event are enriched in multiple signal pathways of metabolism and transport in female placenta. (F) Heatmap showing the profile of splicing factors in male and female placenta with PFHxS expose compare to control.  $N = 6$  in each group, including 3 males and 3 females.  $P$  values were determined by hypergeometric test (D and E). Data in (A-F) are also presented in **Excel Table S21**. Notes: AS, alternative splicing; GO, Gene Ontology; PFHxS, perfluorohexane sulfonate.

**A****C****B****D****E**

**F**



**Figure S5.** AS analyses in multiple signal pathways of placental development after PFHxS exposure. (A) KEGG pathway interaction analysis showing the relationship between signal pathways in RI gene set. (B) KEGG pathway interaction analysis showing the relationship between signal pathways in A3SS gene set. (C) KEGG pathway interaction analysis showing the relationship between signal pathways in A5SS gene set. (D) KEGG pathway interaction analysis showing the relationship between signal pathways in SE gene set. (E) KEGG pathway interaction analysis showing the relationship between signal pathways in MXE gene set. (F) Reactome analysis showing the genes with AS event are enriched in multiple signal pathways related to placental development.  $N = 6$  in each group, including 3 males and 3 females.  $P$  values were determined by hypergeometric test (F). Data in (A-F) are also presented in **Excel Table S22**. Notes: A3SS, alternative 3' splice site; A5SS, alternative 5' splice site; AS, alternative splicing; KEGG, Kyoto Encyclopedia of Genes and Genomes; MXE, exclusion exon; SE skipped exon; PFHxS, perfluorohexane sulfonate; RI, retained intron.



**Figure S6.** Evaluation of placental amino acid transport in the high-dose group and control in both genders. **(A)** The quantitative analysis of mRNA of 12 amino acid transporters in placenta ( $N = 6$ , mean  $\pm$  SEM). **(B)** The profile of 29 amino acids and 44 amides in male placenta ( $N = 5$ ). **(C)** The profile of 29 amino acids and 44 amides in female placenta ( $N = 5$ ).  $P$  values were determined by Student's  $t$  test **(A)**,  $*P < 0.05$ . Data in **(A-C)** are also presented in **Excel Table S23**. Note: PFHxS, perfluorohexane sulfonate; SEM, Standard Error of Mean.